

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No 0704-0188

It is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and reviewing the collection of information, Send comments regarding this burden estimate or any other aspect of reducing this burden, to Washington Headquarters Service, Directorate for Information Operations and Reports, 1215 Jefferson Avenue, S.W., Washington, D.C. 20543, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

## AD-A230 125

Report Date.  
1990

3. Report Type and Dates Covered.  
Abstract

### 5. Funding Numbers.

Program Element No 63207N

Project No X2008

Task No

Accession No DN259001

Evaluation of a Simple Data Assimilation System for Gulf Stream Forecasting

### 6. Author(s).

R. C. Rhodes and T. J. Bennett, Jr.

### 7. Performing Organization Name(s) and Address(es).

Ocean Science Directorate  
Naval Oceanographic and Atmospheric Research Laboratory  
Stennis Space Center, MS 39529-5004

### 8. Performing Organization Report Number.

AB 90:322:110

### 9. Sponsoring/Monitoring Agency Name(s) and Address(es).

Space and Naval Warfare Systems Command  
PDW-141  
Washington, D.C. 20363-5100

### 10. Sponsoring/Monitoring Agency Report Number.

AB 90:322:110

### 11. Supplementary Notes.

EOS

### 12a. Distribution/Availability Statement.

Approved for public release; distribution is unlimited.

### 12b. Distribution Code.

### 13. Abstract (Maximum 200 words).

A Gulf Stream forecast system based on the NOARL two-layer, primitive equation circulation model is being developed and transitioned to operational, U.S. Navy use by the Data Assimilation Research and Transition (DART) team at NOARL and NAVOCEANO. The model is initialized using a dynamic height field obtained by applying the Optimum Thermal Interpolation System (OTIS) feature models to a map of front and ring positions.

The results of experiments designed to evaluate the system in an operationally-realistic mode are presented here. In one set of experiments, observed cloud cover (data gaps) is superimposed on a sequence of frontal paths derived using virtually complete data coverage. In the other set of experiments, operational front and ring maps, rather than research-grade maps, are used for initialization. In both sets of experiments, a rudimentary data assimilation system is established by using a previous forecast to fill in long data gaps in the frontal path.

Both model forecasts, which are initialized from a composite which assume the frontal path does not change in time, are compared to the verification paths. Statistical results indicate that the system's forecast skill is better than that of persistence. In addition, anecdotal results demonstrate the value of the system.

### 14. Subject Terms.

(U) Ocean Models, (U) Data Assimilation

### 15. Number of Pages.

1

### 16. Price Code.

### 20. Limitation of Abstract.

SAR

### 17. Security Classification of Report.

Unclassified


### 18. Security Classification of This Page.

Unclassified

### 19. Security Classification of Abstract.

Unclassified

DTIC  
ELECTE  
DEC 20 1990  
S B D



A-1,211